


Figure 1 of 26



BEST AVAILABLE COPY

Figure 2 of 26




Figure 3 of 26

Oligosaccharide	m/z	MRM pair	P2 fraction ^b	GAG ^c
IdoA- GalNAc4SQ-UA- GalNAc- UA (+SQ)	720	720/269	#31 and #32	DS
IdoA- GalNAc4SQ-UA- GalNAc (+SQ)	632	632/298	#34 and #35	DS
IdoA- GalNAc4SQ-UA	982	982/269	#38 and #39	DS
	490	490/173		
IdoA-HN-UA (+SQ)	940	940/269	#37	HS
IdoA-HN-UA (+2SQ)	1020	1020/940	#37	HS
	509	509/422		
IdoA- HNAc (+SQ)	806	806/295	#40 and #41	HS/DS

^a IdoA, iduronic acid; GalNAc, N-acetylgalactosamine; GalNAc4SQ, N-acetylgalactosamine

4 - sulfate; UA, uronic acid; HN, hexosamine; HNAc, N-acetylhexosamine.

^b fractions from the Bio- Gel P2 column (see Figure 23).

^c proposed source of oligosaccharide (DS, dermatan sulfate; HS, heparan sulfate).

Figure 4 of 26



Figure 5 of 26






Figure 6 of 26

A



B




Figure 7 of 26




Figure 8 of 26

Relative oligosaccharide levels in control^a urine

Oligosaccharide ^b	Q1/Q3 ^c	Average value ^d	Std. Dev.
UA	523/173	0.1480	0.1119
HNAC	550/173	0.0036	0.0028
HNACS	630/256	0.0016	0.0007
HNACS2	710/256	0.0030	0.0018
HNAC-UA	726/331	0.0155	0.0196
(HNAC, UA)	726/173	0.0063	0.0072
(HNS, UA)	764/173	0.0736	0.0295
HNS-UA	764/331	0.1148	0.1041
(HNAC, UA, S)	806/173	0.0003	0.0003
UA-HNACS	806/295	0.1941	0.1198
HNACS-UA	806/331	0.0999	0.0882
UA-HN-UA (S)	940/269	0.0294	0.0106
UA-HNAC-UA (S)	982/269	0.0111	0.0120
UA-HNAC-UA (S)	490/173	0.0045	0.0029
UA-HNAC-UA (S)	490/476	0.0305	0.0161
UA-HN-UA (S2)	1020/269	0.0026	0.0013
UA-HN-UA (S2)	1020/940	0.0018	0.0015
UA-HN-UA (S2)	509/422	0.0011	0.0018
HNAC-UA-HNAC (S2)	544/173	0.0507	0.0345
HNAC-UA-HNAC-UA (S)	1185/931	0.0099	0.0080
HNAC-UA-HNAC-UA (S)	591/173	0.2024	0.1818
HNAC-UA-HNAC-UA (S2)	632/298	0.0556	0.0358
UA-HNAC-UA-HNAC-UA (S2)	720/269	0.0024	0.0024
HNAC-UA-HNAC-UA-HNAC	734/173	0.0044	0.0027
HNAC-UA-HNAC-UA-HNAC (S3)	515/173	0.0291	0.0112
Unknown	528/448	0.0195	0.0082
Hexasaccharide	764/256	0.0007	0.0006

^a Controls = 26^b UA=uronic acid, HN=hexosamine, HNAC=N-acetylhexosamine, S=sulphate, structures in brackets indicate unknown sequence.^c m/z values used for the MRM analysis of selected oligosaccharides.^d values refer to the ratio values of the oligosaccharide signals to the internal standards GlcNAc6S (d3) (used for the monosaccharides) and the chondroitin disaccharide α -ΔUA-[1-3]-GalNAc-6S (used for disaccharides and larger species).

Figure 9 of 26

Mann Whitney U values for oligosaccharide analytes in MPS subgroups^d compared to controls^e

Oligosaccharide ^a	MPS I 15	MPS II 13	MPS IIIA 12	MPS IIIB 7	MPS IIIC 2	MPS IIID 2	MPS IVA 6	MPS VI 10	MS 1
UA	184	156.5	136	75	19.5 ^c	24.5	70	83	2
HNAC	128	123.5	99	79	40	26	53	106	5
HNACs	73 ^b	101 ^c	67 ^b	53	27	0 ^c	0 ^b	0 ^b	5.5
HNACs2	85 ^b	127	60 ^b	58	22	14	30 ^c	7 ^b	4
HNAC-UA	170.5	134.5	117	43 ^c	29	21	51	55 ^b	3
(HNAC, UA)	170	123	86 ^c	77	23.5	10	39	62 ^c	5
(HNS, UA)	5 ^b	4 ^b	31 ^b	49.5	25	1 ^c	32 ^c	104	0
HNS-UA	114.5 ^c	17 ^b	0 ^b	51	44.5	1 ^c	42	75.5	0
(HNAC, UA, S)	150	71.5 ^b	81.5 ^c	26 ^b	50	6	54	86	5
UA-HNACs	0 ^b	0 ^b	120.5	80	42	2 ^c	67	115	8
HNACs-UA	194	152	154	76	38	0 ^c	1 ^b	4 ^b	3
UA-HN-UA (S)	15.5 ^b	140	139.5	91	35.5	9	63	77	0
UA-HNAC-UA (S)	0 ^b	53 ^b	146	88	35	21	70.5	90	7
UA-HNAC-UA (S)	75.5 ^b	153	140	65	45	5	38	108	7
UA-HNAC-UA (S)	0 ^b	8 ^b	140	51	36	23	67.5	128	5
UA-HN-UA (S2)	169	52 ^b	66 ^b	40 ^c	7 ^b	21	37 ^c	48 ^b	9
UA-HN-UA (S2)	86.5 ^b	134	113.5	81	48	22	71	115	1
UA-HN-UA (S2)	24 ^b	136	69 ^b	76	46	22	75	96	0
HNAC-UA-HNAC (S2)	109 ^c	78 ^b	133.5	78	46	10	15.5 ^b	62 ^c	5
HNAC-UA-HNAC-UA (S)	0 ^b	134	64 ^b	0 ^b	22	3 ^c	10 ^b	29 ^b	0
HNAC-UA-HNAC-UA (S)	61 ^b	68 ^b	128	34 ^c	37	11	56	118.5	13
HNAC-UA-HNAC-UA (S2)	5 ^b	130	31 ^b	5 ^b	28	0 ^c	10.5 ^b	0 ^b	0
UA-HNAC-UA-HNAC-UA (S2)	61.5 ^b	131	101.5	71.5	31	8	70	122.5	2
HNAC-UA-HNAC-UA-HNAC	188	76.5 ^b	90 ^c	63	27	17	50	108	4
HNAC-UA-HNAC-UA-HNAC (S3)	167	55.5 ^b	125	44.5 ^c	50	17	68	105	11
Unknown (S28/448)	60.5 ^b	54 ^b	43 ^b	57	24	8.5	41	110	2
Hexasaccharide (764/256)	0 ^b	0 ^b	13 ^b	1 ^b	3 ^b	0 ^c	2 ^b	8 ^b	0

^a UA=uronic acid, HN=hexosamine, HNAC=N-acetylhexosamine, S=sulphate, structures in brackets indicate unknown sequence.^b p<0.01^c p<0.5^d MPS types are shown with the number of patient samples below

Figure 10 of 26





Figure 11 of 26



12/26

Figure 12 of 26




Figure 13 of 26

Cell line	Disorder	UA-HN-UA(Sx2) ¹
SF 5344	Control	0.034
SF 5248	Control	0.030
SF 2662	MPS I (intermediate)	0.090
SF 5048	MPS I (severe)	0.379
SF 538	MPS I (severe)	0.387

¹ relative level of oligosaccharide/mg cell protein
UA = uronic acid; HN = hexosamine

Figure 14 of 26

Genotype	632/298 (MRM)	982/269 (MRM)	940/269 (MRM)	1020/269 (MRM)	509/422 (MRM)	806/295 (MRM)
W402X _{hom}	30.8	93.6	14.0	4.8	20.0	111.8
W402X, Q70X	5.4	10.9	1.0	3.3	1.0	51.4
W180X, c134-145del	5.6	5.7	0.55	0.69	1.6	25.2
P533R, W402X	4.4	5.7	0.67	0.32	1.1	40.9
R89Q, 1060+2t>c	3.4	2.7	0.13	0.33	0.89	11.3
R89Q, 1060+2t>c	5.4	2.2	0.19	0.19	1.0	10.5
L346R, c704ins5	2.7	1.5	0.21	0.18	0.68	9.1
Control range (n=26)	0.07-0.9	0.07-0.6	0.02-0.2	0-4.0	0-0.6	0.03-0.5

Figure 15 of 26

Oligosaccharide ¹ MRM pairs	Pre-BMT	Post-BMT (3 months)	Post-BMT (8 months)	Control Avg (n=8)	Control Std Dev
UA-HN-UA(Sx2)	509/422	2.31	0.63	0.59	0.03
HNAC(S)	630/256	192	165	139	52
UA-HNAC(S)	806/295	13.93	8.39	2.12	0.02
UA-HN-UA(S)	940/269	0.49	0.21	0.19	0.02
UA-HNAC- UA(S)	982/269	2.58	1.86	1.05	0.04
UA-HNAC-UA- HNAC(Sx2)	632/298	9.34	4.51	2.57	0.16

¹UA = uronic acid; HN = hexosamine; S = sulphate; HNAC = N-acetylhexosamine

Figure 16 of 26




Figure 17 of 26




Figure 18 of 26




Figure 19 of 26




Figure 20 of 26




Figure 21 of 26




Figure 22 of 26




Figure 23 of 26




Figure 24 of 26




Figure 25 of 26





Figure 26 of 26



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.